Desired Performance	Target Measures	Baseline	Ballad	Health	
P A			FY23	FYTD24	
•	PSI 3 Pressure Ulcer Rate	1.07	0.10	0.35	
•	PSI 6 latrogenic Pneumothorax Rate	0.25	0.06	0.02	
•	PSI 8a In Hospital Fall with Hip Fracture Rate	0.06	0.09	0.02	
•	PSI 8b In Hospital Fall with Other Fracture Rate			0.29	
•	PSI 9 Perioperative Hemorrhage or Hematoma Rate	1.59	1.46	1.56	
•	PSI 10 Postoperative Acute Kidney Injury Requiring Dialysis Rate	0.76	2.40	1.80	
•	PSI 11 Postoperative Respiratory Failure Rate	9.24	5.30	4.09	
•	PSI 12 Perioperative Pulmonary Embolism or Deep Vein Thrombosis Rate	3.31	2.51	2.88	
•	PSI 13 Postoperative Sepsis Rate	3.58	3.17	4.77	
•	PSI 14 Postoperative Wound Dehiscence Rate	0.83	2.14	1.92	
	PSI 15 Unrecognized Abdominopelvic Accidental Puncture/Laceration Rate	1.18	0.62	0.00	
•	SMB: Sepsis Management Bundle	56.9%	59.2%	60.9%	
₩	CLABSI	0.711	1.037	0.695	
•	CAUTI	0.558	0.729	0.724	
•	SSI COLON Surgical Site Infection	2.13	2.94	3.13	
•	SSI HYST Surgical Site Infection	0.71	1.51	2.30	
•	MRSA	0.047	0.080	0.047	
•	CDIFF	0.671	0.182	0.115	
<b>1</b>	HCOMP1A P Patients who reported that their nurses "Always" communicated well	84.6	77.2	79.5	
•	HCOMP2A P Patients who reported that their doctors "Always" communicated well	83.3	78.3	78.6	
•	HCOMP5A P Patients who reported that staff "Always" explained about medicines before giving it to them	71.0	61.9	65.5	
•	HCOMP6Y P Patients who reported that YES, they were given information about what to do during their recovery at home	87.7	86.0	85.8	
•	HCOMP7SA Patients who "Strongly Agree" they understood their care when they left the hospital	54.6	47.9	49.6	
₩.	READM30HF Heart Failure 30day readmissions rate	27.7%	24.2%	24.2%	
	READM30PN Pneumonia 30day readmission rate	17.9%	18.4%	18.0%	
	Sepsis In House Mortality	10.7%	9.8%	8.9%	
	MORT30HF Heart failure 30day mortality rate	3.0%	3.6%	3.1%	
•	MORT30PN Pneumonia 30day mortality rate	5.0%	5.0%	4.9%	
•	Left without being seen	0.83%	1.37%	0.90%	
•	Median Time from ED Arrival to Departure for Outpatients (18b)	126.00	160.8	152.2	
•	Median Time from ED Arrival to Transport for Admitted Patients (ED1)	223.00	603.3	441.6	

Desired Performance	Target Measures	Baseline		ity Medical nter
Pe			FY23	FYTD24
•	PSI 3 Pressure Ulcer Rate	1.07	0.14	0.37
•	PSI 6 latrogenic Pneumothorax Rate	0.25	0.00	0.00
•	PSI 8a In Hospital Fall with Hip Fracture Rate	0.06	0.05	0.00
•	PSI 8b In Hospital Fall with Other Fracture Rate			0.28
•	PSI 9 Perioperative Hemorrhage or Hematoma Rate	1.59	1.38	0.90
	PSI 10 Postoperative Acute Kidney Injury Requiring Dialysis Rate	0.76	2.25	1.81
•	PSI 11 Postoperative Respiratory Failure Rate	9.24	6.16	1.94
•	PSI 12 Perioperative Pulmonary Embolism or Deep Vein Thrombosis Rate	3.31	2.23	2.30
•	PSI 13 Postoperative Sepsis Rate	3.58	2.95	6.55
•	PSI 14 Postoperative Wound Dehiscence Rate	0.83	2.59	5.38
+	PSI 15 Unrecognized Abdominopelvic Accidental Puncture/Laceration	1.18	0.74	0.00
•	SMB: Sepsis Management Bundle	56.9%	51.6%	29.7%
•	CLABSI	0.711	1.558	0.769
•	CAUTI	0.558	1.055	1.587
•	SSI COLON Surgical Site Infection	2.13	4.11	7.23
•	SSI HYST Surgical Site Infection	0.71	0.00	0.00
•	MRSA	0.047	0.093	0.054
<b>+</b>	CDIFF	0.671	0.122	0.121
<b>†</b>	HCOMP1A P Patients who reported that their nurses "Always" communicated well	84.6	72.2	70.6
•	HCOMP2A P Patients who reported that their doctors "Always" communicated well	83.3	72.3	71.9
•	HCOMP5A P Patients who reported that staff "Always" explained about medicines before giving it to them	71.0	57.6	55.7
•	HCOMP6Y P Patients who reported that YES, they were given information about what to do during their recovery at home	87.7	82.4	82.6
•	HCOMP7SA Patients who "Strongly Agree" they understood their care when they left the hospital	54.6	44.0	41.3
•	READM30HF Heart Failure 30day readmissions rate	27.7%	26.5%	26.8%
•	READM30PN Pneumonia 30day readmission rate	17.9%	18.9%	18.1%
•	Sepsis In House Mortality	10.7%	18.1%	13.7%
•	MORT30HF Heart failure 30day mortality rate	3.0%	3.0%	4.1%
•	MORT30PN Pneumonia 30day mortality rate	5.0%	8.5%	6.6%
•	Left without being seen	0.83%	0.96%	0.84%
•	Median Time from ED Arrival to Departure for Outpatients (18b)	126.00	196.1	207.1
•	Median Time from ED Arrival to Transport for Admitted Patients (ED1)	223.00	506.5	613.1

Desired Performance	Target Measures	Baseline		n Valley I Center
Per			FY23	FYTD24
•	PSI 3 Pressure Ulcer Rate	1.07	0.11	0.30
•	PSI 6 latrogenic Pneumothorax Rate	0.25	0.15	0.00
•	PSI 8a In Hospital Fall with Hip Fracture Rate	0.06	0.14	0.00
•	PSI 8b In Hospital Fall with Other Fracture Rate			0.21
•	PSI 9 Perioperative Hemorrhage or Hematoma Rate	1.59	1.16	1.04
	PSI 10 Postoperative Acute Kidney Injury Requiring Dialysis Rate	0.76	4.04	4.12
•	PSI 11 Postoperative Respiratory Failure Rate	9.24	6.65	6.49
•	PSI 12 Perioperative Pulmonary Embolism or Deep Vein Thrombosis Rate	3.31	3.10	3.57
•	PSI 13 Postoperative Sepsis Rate	3.58	3.06	4.26
•	PSI 14 Postoperative Wound Dehiscence Rate	0.83	0.00	0.00
•	PSI 15 Unrecognized Abdominopelvic Accidental Puncture/Laceration	1.18	1.05	0.00
•	SMB: Sepsis Management Bundle	56.9%	50.8%	41.2%
	CLABSI	0.711	1.004	1.105
•	CAUTI	0.558	0.755	0.970
•	SSI COLON Surgical Site Infection	2.13	2.92	6.78
•	SSI HYST Surgical Site Infection	0.71	4.23	0.00
•	MRSA	0.047	0.097	0.074
•	CDIFF	0.671	0.304	0.129
<b>1</b>	HCOMP1A P Patients who reported that their nurses "Always" communicated well	84.6	70.0	72.7
<b>1</b>	HCOMP2A P Patients who reported that their doctors "Always" communicated well	83.3	75.0	76.2
<b>1</b>	HCOMP5A P Patients who reported that staff "Always" explained about medicines before giving it to them	71.0	52.3	54.7
<b>+</b>	HCOMP6Y P Patients who reported that YES, they were given information about what to do during their recovery at home	87.7	86.4	86.2
•	HCOMP7SA Patients who "Strongly Agree" they understood their care when they left the hospital	54.6	44.2	45.2
•	READM30HF Heart Failure 30day readmissions rate	27.7%	21.5%	23.2%
•	READM30PN Pneumonia 30day readmission rate	17.9%	17.9%	17.5%
•	Sepsis In House Mortality	10.7%	13.1%	13.4%
<b>+</b>	MORT30HF Heart failure 30day mortality rate	3.0%	4.6%	3.1%
•	MORT30PN Pneumonia 30day mortality rate	5.0%	6.9%	6.3%
•	Left without being seen	0.83%	0.63%	0.46%
•	Median Time from ED Arrival to Departure for Outpatients (18b)	126.00	215.8	207.9
•	Median Time from ED Arrival to Transport for Admitted Patients (ED1)	223.00	943.7	528.1

Desired Performance	Target Measures	Baseline		Regional I Center
Pe			FY23	FYTD24
•	PSI 3 Pressure Ulcer Rate	1.07	0.11	0.00
•	PSI 6 latrogenic Pneumothorax Rate	0.25	0.08	0.00
•	PSI 8a In Hospital Fall with Hip Fracture Rate	0.06	0.16	0.00
•	PSI 8b In Hospital Fall with Other Fracture Rate			0.45
•	PSI 9 Perioperative Hemorrhage or Hematoma Rate	1.59	2.65	0.72
•	PSI 10 Postoperative Acute Kidney Injury Requiring Dialysis Rate	0.76	2.52	0.00
•	PSI 11 Postoperative Respiratory Failure Rate	9.24	2.70	8.51
•	PSI 12 Perioperative Pulmonary Embolism or Deep Vein Thrombosis Rate	3.31	0.81	3.34
•	PSI 13 Postoperative Sepsis Rate	3.58	4.13	4.28
•	PSI 14 Postoperative Wound Dehiscence Rate	0.83	1.90	0.00
	PSI 15 Unrecognized Abdominopelvic Accidental Puncture/Laceration Rate	1.18	0.91	0.00
•	SMB: Sepsis Management Bundle	56.9%	47.8%	58.7%
-	CLABSI	0.711	1.029	0.173
•	CAUTI	0.558	0.838	0.633
•	SSI COLON Surgical Site Infection	2.13	2.84	1.05
•	SSI HYST Surgical Site Infection	0.71	0.00	0.00
•	MRSA	0.047	0.121	0.062
<b>+</b>	CDIFF	0.671	0.188	0.187
<b>+</b>	HCOMP1A P Patients who reported that their nurses "Always" communicated well	84.6	71.1	74.7
•	HCOMP2A P Patients who reported that their doctors "Always" communicated well	83.3	75.3	75.0
•	HCOMP5A P Patients who reported that staff "Always" explained about medicines before giving it to them	71.0	58.5	59.1
•	HCOMP6Y P Patients who reported that YES, they were given information about what to do during their recovery at home	87.7	86.5	85.7
•	HCOMP7SA Patients who "Strongly Agree" they understood their care when they left the hospital	54.6	47.2	47.7
•	READM30HF Heart Failure 30day readmissions rate	27.7%	22.4%	24.7%
•	READM30PN Pneumonia 30day readmission rate	17.9%	20.6%	21.2%
•	Sepsis In House Mortality	10.7%	11.4%	9.6%
•	MORT30HF Heart failure 30day mortality rate	3.0%	3.9%	2.7%
•	MORT30PN Pneumonia 30day mortality rate	5.0%	5.5%	6.3%
•	Left without being seen	0.83%	2.15%	1.29%
•	Median Time from ED Arrival to Departure for Outpatients (18b)	126.00	217.5	204.3
•	Median Time from ED Arrival to Transport for Admitted Patients (ED1)	223.00	617.2	381.1

Desired Performance	Target Measures	Baseline		Memorial pital
Pe			FY23	FYTD24
•	PSI 3 Pressure Ulcer Rate	1.07	0.00	1.27
₩	PSI 6 latrogenic Pneumothorax Rate	0.25	0.00	0.28
<b>+</b>	PSI 8a In Hospital Fall with Hip Fracture Rate	0.06	0.19	0.00
•	PSI 8b In Hospital Fall with Other Fracture Rate			0.27
•	PSI 9 Perioperative Hemorrhage or Hematoma Rate	1.59	0.00	3.59
	PSI 10 Postoperative Acute Kidney Injury Requiring Dialysis Rate	0.76	0.00	0.00
•	PSI 11 Postoperative Respiratory Failure Rate	9.24	0.00	0.00
	PSI 12 Perioperative Pulmonary Embolism or Deep Vein Thrombosis Rate	3.31	5.27	5.05
•	PSI 13 Postoperative Sepsis Rate	3.58	0.00	0.00
•	PSI 14 Postoperative Wound Dehiscence Rate	0.83	0.00	0.00
+	PSI 15 Unrecognized Abdominopelvic Accidental Puncture/Laceration Rate	1.18	0.00	0.00
•	SMB: Sepsis Management Bundle	56.9%	58.6%	61.5%
•	CLABSI	0.711	0.366	1.007
	CAUTI	0.558	1.162	0.000
•	SSI COLON Surgical Site Infection	2.13	3.49	0.00
•	SSI HYST Surgical Site Infection	0.71	0.00	0.00
•	MRSA	0.047	0.035	0.000
<b>+</b>	CDIFF	0.671	0.107	0.053
•	HCOMP1A P Patients who reported that their nurses "Always" communicated well	84.6	72.6	77.1
<b>1</b>	HCOMP2A P Patients who reported that their doctors "Always" communicated well	83.3	76.3	81.1
<b>1</b>	HCOMP5A P Patients who reported that staff "Always" explained about medicines before giving it to them	71.0	54.6	54.8
•	HCOMP6Y P Patients who reported that YES, they were given information about what to do during their recovery at home	87.7	85.5	87.8
<b>1</b>	HCOMP7SA Patients who "Strongly Agree" they understood their care when they left the hospital	54.6	43.8	48.5
•	READM30HF Heart Failure 30day readmissions rate	27.7%	24.7%	25.9%
•	READM30PN Pneumonia 30day readmission rate	17.9%	18.9%	20.2%
•	Sepsis In House Mortality	10.7%	8.6%	9.9%
•	MORT30HF Heart failure 30day mortality rate	3.0%	3.1%	1.9%
•	MORT30PN Pneumonia 30day mortality rate	5.0%	5.0%	5.7%
•	Left without being seen	0.83%	1.59%	1.65%
•	Median Time from ED Arrival to Departure for Outpatients (18b)	126.00	213.5	234.5
•	Median Time from ED Arrival to Transport for Admitted Patients (ED1)	223.00	507.8	434.8

Desired Performance	Target Measures	Baseline		Woods ty Hospital
D Perf			FY23	FYTD24
•	PSI 3 Pressure Ulcer Rate	1.07	0.00	0.49
•	PSI 6 latrogenic Pneumothorax Rate	0.25	0.00	0.00
•	PSI 8a In Hospital Fall with Hip Fracture Rate	0.06	0.00	0.28
•	PSI 8b In Hospital Fall with Other Fracture Rate	0.00	0.00	0.55
•	PSI 9 Perioperative Hemorrhage or Hematoma Rate	1.59	2.17	3.34
	PSI 10 Postoperative Acute Kidney Injury Requiring Dialysis Rate	0.76	0.00	0.00
•	PSI 11 Postoperative Respiratory Failure Rate	9.24	4.89	0.00
	PSI 12 Perioperative Pulmonary Embolism or Deep Vein Thrombosis		4.03	0.00
	Rate	3.31	1.02	0.00
•	PSI 13 Postoperative Sepsis Rate	3.58	2.55	3.40
•	PSI 14 Postoperative Wound Dehiscence Rate	0.83	2.24	0.00
•	PSI 15 Unrecognized Abdominopelvic Accidental Puncture/Laceration	1.18	0.00	0.00
<b>1</b>	SMB: Sepsis Management Bundle	56.9%	64.4%	68.2%
₩	CLABSI	0.711	0.000	1.374
•	CAUTI	0.558	0.000	0.000
•	SSI COLON Surgical Site Infection	2.13	0.48	3.57
•	SSI HYST Surgical Site Infection	0.71	0.00	0.00
•	MRSA	0.047	0.000	0.055
•	CDIFF	0.671	0.080	0.061
•	HCOMP1A P Patients who reported that their nurses "Always" communicated well	84.6	81.4	82.3
•	HCOMP2A P Patients who reported that their doctors "Always" communicated well	83.3	82.3	79.8
•	HCOMP5A P Patients who reported that staff "Always" explained about medicines before giving it to them	71.0	67.6	67.1
•	HCOMP6Y P Patients who reported that YES, they were given information about what to do during their recovery at home	87.7	86.7	88.3
•	HCOMP7SA Patients who "Strongly Agree" they understood their care when they left the hospital	54.6	54.1	55.6
	READM30HF Heart Failure 30day readmissions rate	27.7%	25.2%	21.0%
•	READM30PN Pneumonia 30day readmission rate	17.9%	18.4%	13.5%
•	Sepsis In House Mortality	10.7%	3.4%	3.0%
•	MORT30HF Heart failure 30day mortality rate	3.0%	2.7%	3.0%
•	MORT30PN Pneumonia 30day mortality rate	5.0%	3.3%	2.9%
•	Left without being seen	0.83%	1.72%	1.25%
•	Median Time from ED Arrival to Departure for Outpatients (18b)	126.00	199.8	190.6
•	Median Time from ED Arrival to Transport for Admitted Patients (ED1)	223.00	777.8	702.5

ed ance				n Path
Desired Performance	Target Measures	Baseline	Communi	ty Hospital
ď			FY23	FYTD24
	PSI 3 Pressure Ulcer Rate	1.07	0.00	0.00
•	PSI 6 latrogenic Pneumothorax Rate	0.25	0.00	0.00
•	PSI 8a In Hospital Fall with Hip Fracture Rate	0.06	0.00	0.00
•	PSI 8b In Hospital Fall with Other Fracture Rate			0.00
	PSI 9 Perioperative Hemorrhage or Hematoma Rate	1.59	0.00	0.00
•	PSI 10 Postoperative Acute Kidney Injury Requiring Dialysis Rate	0.76	0.00	0.00
•	PSI 11 Postoperative Respiratory Failure Rate	9.24	0.00	0.00
•	PSI 12 Perioperative Pulmonary Embolism or Deep Vein Thrombosis Rate	3.31	0.00	7.25
•	PSI 13 Postoperative Sepsis Rate	3.58	10.00	0.00
•	PSI 14 Postoperative Wound Dehiscence Rate	0.83	0.00	13.16
•	PSI 15 Unrecognized Abdominopelvic Accidental Puncture/Laceration Rate	1.18	2.70	0.00
•	SMB: Sepsis Management Bundle	56.9%	67.8%	81.1%
<del>-</del>	CLABSI	0.711	0.000	0.000
•	CAUTI	0.558	0.000	0.000
•	SSI COLON Surgical Site Infection	2.13	4.48	0.00
•	SSI HYST Surgical Site Infection	0.71	0.00	0.00
	MRSA	0.047	0.199	0.000
	CDIFF	0.671	0.381	0.000
•	HCOMP1A P Patients who reported that their nurses "Always" communicated well	84.6	80.7	78.5
•	HCOMP2A P Patients who reported that their doctors "Always" communicated well	83.3	81.6	79.9
•	HCOMP5A P Patients who reported that staff "Always" explained about medicines before giving it to them	71.0	66.3	66.8
•	HCOMP6Y P Patients who reported that YES, they were given information about what to do during their recovery at home	87.7	84.9	85.5
•	HCOMP7SA Patients who "Strongly Agree" they understood their care when they left the hospital	54.6	51.9	49.5
•	READM30HF Heart Failure 30day readmissions rate	27.7%	24.4%	10.5%
•	READM30PN Pneumonia 30day readmission rate	17.9%	16.8%	15.7%
•	Sepsis In House Mortality	10.7%	0.2%	0.4%
•	MORT30HF Heart failure 30day mortality rate	3.0%	2.4%	0.0%
•	MORT30PN Pneumonia 30day mortality rate	5.0%	0.0%	0.0%
•	Left without being seen	0.83%	1.03%	0.33%
•	Median Time from ED Arrival to Departure for Outpatients (18b)	126.00	170.9	149.0
•	Median Time from ED Arrival to Transport for Admitted Patients (ED1)	223.00	789.1	420.2

Desired Performance	Target Measures	Baseline		ommunity pital
Pe			FY23	FYTD24
+	PSI 3 Pressure Ulcer Rate	1.07	0.00	0.91
<b>+</b>	PSI 6 latrogenic Pneumothorax Rate	0.25	0.00	0.00
•	PSI 8a In Hospital Fall with Hip Fracture Rate	0.06	0.00	0.00
<b>+</b>	PSI 8b In Hospital Fall with Other Fracture Rate			0.00
•	PSI 9 Perioperative Hemorrhage or Hematoma Rate	1.59	0.00	0.00
•	PSI 10 Postoperative Acute Kidney Injury Requiring Dialysis Rate	0.76	0.00	0.00
•	PSI 11 Postoperative Respiratory Failure Rate	9.24	0.00	0.00
•	PSI 12 Perioperative Pulmonary Embolism or Deep Vein Thrombosis Rate	3.31	2.54	4.05
•	PSI 13 Postoperative Sepsis Rate	3.58	0.00	0.00
•	PSI 14 Postoperative Wound Dehiscence Rate	0.83	5.46	0.00
•	PSI 15 Unrecognized Abdominopelvic Accidental Puncture/Laceration Rate	1.18	0.00	0.00
<b>1</b>	SMB: Sepsis Management Bundle	56.9%	57.1%	66.7%
<u> </u>	CLABSI	0.711	1.918	0.000
•	CAUTI	0.558	0.000	0.000
•	SSI COLON Surgical Site Infection	2.13	10.42	4.17
•	SSI HYST Surgical Site Infection	0.71	0.00	0.00
•	MRSA	0.047	0.000	0.000
<b>+</b>	CDIFF	0.671	0.087	0.130
<b>1</b>	HCOMP1A P Patients who reported that their nurses "Always" communicated well	84.6	70.3	76.7
<b>1</b>	HCOMP2A P Patients who reported that their doctors "Always" communicated well	83.3	71.6	75.9
<b>1</b>	HCOMP5A P Patients who reported that staff "Always" explained about medicines before giving it to them	71.0	50.5	58.8
•	HCOMP6Y P Patients who reported that YES, they were given information about what to do during their recovery at home	87.7	79.6	81.6
<b>1</b>	HCOMP7SA Patients who "Strongly Agree" they understood their care when they left the hospital	54.6	37.2	46.4
•	READM30HF Heart Failure 30day readmissions rate	27.7%	25.4%	21.2%
•	READM30PN Pneumonia 30day readmission rate	17.9%	17.4%	19.5%
•	Sepsis In House Mortality	10.7%	5.6%	3.8%
•	MORT30HF Heart failure 30day mortality rate	3.0%	3.3%	3.0%
•	MORT30PN Pneumonia 30day mortality rate	5.0%	3.3%	1.9%
•	Left without being seen	0.83%	1.83%	0.87%
•	Median Time from ED Arrival to Departure for Outpatients (18b)	126.00	160.4	146.4
•	Median Time from ED Arrival to Transport for Admitted Patients (ED1)	223.00	879.2	696.8

ed ance				neville
Desired Performance	Target Measures	Baseline	Communi	ty Hospital
ď			FY23	FYTD24
•	PSI 3 Pressure Ulcer Rate	1.07	0.00	0.54
•	PSI 6 latrogenic Pneumothorax Rate	0.25	0.25	0.00
•	PSI 8a In Hospital Fall with Hip Fracture Rate	0.06	0.00	0.00
•	PSI 8b In Hospital Fall with Other Fracture Rate			0.68
•	PSI 9 Perioperative Hemorrhage or Hematoma Rate	1.59	2.76	5.60
•	PSI 10 Postoperative Acute Kidney Injury Requiring Dialysis Rate	0.76	0.00	0.00
•	PSI 11 Postoperative Respiratory Failure Rate	9.24	14.08	8.33
•	PSI 12 Perioperative Pulmonary Embolism or Deep Vein Thrombosis Rate	3.31	3.85	3.50
•	PSI 13 Postoperative Sepsis Rate	3.58	7.87	9.52
•	PSI 14 Postoperative Wound Dehiscence Rate	0.83	3.51	0.00
•	PSI 15 Unrecognized Abdominopelvic Accidental Puncture/Laceration Rate	1.18	0.00	0.00
•	SMB: Sepsis Management Bundle	56.9%	59.5%	50.0%
<del>-</del>	CLABSI	0.711	1.235	0.755
•	CAUTI	0.558	0.000	0.000
•	SSI COLON Surgical Site Infection	2.13	1.03	1.14
•	SSI HYST Surgical Site Infection	0.71	0.00	
•	MRSA	0.047	0.049	0.000
	CDIFF	0.671	0.255	0.073
•	HCOMP1A P Patients who reported that their nurses "Always" communicated well	84.6	72.0	75.8
•	HCOMP2A P Patients who reported that their doctors "Always" communicated well	83.3	73.8	75.9
•	HCOMP5A P Patients who reported that staff "Always" explained about medicines before giving it to them	71.0	63.2	59.6
•	HCOMP6Y P Patients who reported that YES, they were given information about what to do during their recovery at home	87.7	84.6	87.2
•	HCOMP7SA Patients who "Strongly Agree" they understood their care when they left the hospital	54.6	43.2	46.0
•	READM30HF Heart Failure 30day readmissions rate	27.7%	27.1%	29.7%
•	READM30PN Pneumonia 30day readmission rate	17.9%	17.0%	18.5%
•	Sepsis In House Mortality	10.7%	5.6%	5.5%
•	MORT30HF Heart failure 30day mortality rate	3.0%	2.6%	3.0%
•	MORT30PN Pneumonia 30day mortality rate	5.0%	2.1%	3.6%
•	Left without being seen	0.83%	1.42%	1.10%
•	Median Time from ED Arrival to Departure for Outpatients (18b)	126.00	187.9	169.4
•	Median Time from ED Arrival to Transport for Admitted Patients (ED1)	223.00	772.9	680.2

Desired Performance	Target Measures	Baseline	1	County ty Hospital
Per			FY23	FYTD24
+	PSI 3 Pressure Ulcer Rate	1.07	0.00	0.00
•	PSI 6 latrogenic Pneumothorax Rate	0.25	0.00	0.00
₩	PSI 8a In Hospital Fall with Hip Fracture Rate	0.06	0.00	0.00
•	PSI 8b In Hospital Fall with Other Fracture Rate	0.00	0.00	0.00
•	PSI 9 Perioperative Hemorrhage or Hematoma Rate	1.59	0.00	0.00
	PSI 10 Postoperative Acute Kidney Injury Requiring Dialysis Rate	0.76	0.00	0.00
1	PSI 11 Postoperative Respiratory Failure Rate	9.24	0.00	0.00
,	PSI 12 Perioperative Pulmonary Embolism or Deep Vein Thrombosis		0.00	
	Rate	3.31	13.16	0.00
	PSI 13 Postoperative Sepsis Rate	3.58	0.00	0.00
	PSI 14 Postoperative Wound Dehiscence Rate	0.83	0.00	0.00
•	PSI 15 Unrecognized Abdominopelvic Accidental Puncture/Laceration Rate	1.18	0.00	0.00
•	SMB: Sepsis Management Bundle	56.9%	84.6%	87.2%
-	CLABSI	0.711	0.000	0.000
•	CAUTI	0.558	0.000	0.000
•	SSI COLON Surgical Site Infection	2.13		
•	SSI HYST Surgical Site Infection	0.71		
•	MRSA	0.047	0.000	0.000
•	CDIFF	0.671	0.000	0.000
•	HCOMP1A P Patients who reported that their nurses "Always" communicated well	84.6	77.2	80.5
•	HCOMP2A P Patients who reported that their doctors "Always" communicated well	83.3	78.1	78.6
•	HCOMP5A P Patients who reported that staff "Always" explained about medicines before giving it to them	71.0	65.8	66.4
•	HCOMP6Y P Patients who reported that YES, they were given information about what to do during their recovery at home	87.7	86.2	86.3
•	HCOMP7SA Patients who "Strongly Agree" they understood their care when they left the hospital	54.6	48.8	48.2
•	READM30HF Heart Failure 30day readmissions rate	27.7%	26.7%	7.7%
•	READM30PN Pneumonia 30day readmission rate	17.9%	18.8%	15.0%
•	Sepsis In House Mortality	10.7%	4.2%	4.8%
•	MORT30HF Heart failure 30day mortality rate	3.0%	4.3%	8.3%
+	MORT30PN Pneumonia 30day mortality rate	5.0%	2.5%	2.5%
+	Left without being seen	0.83%	1.89%	1.17%
•	Median Time from ED Arrival to Departure for Outpatients (18b)	126.00	138.2	140.3
•	Median Time from ED Arrival to Transport for Admitted Patients (ED1)	223.00	383.0	593.1

Desired Performance	Target Measures	Baseline	•	e Shoals pital
Pel			FY23	FYTD24
<b>+</b>	PSI 3 Pressure Ulcer Rate	1.07	0.51	0.00
•	PSI 6 latrogenic Pneumothorax Rate	0.25	0.00	0.00
•	PSI 8a In Hospital Fall with Hip Fracture Rate	0.06	0.32	0.00
•	PSI 8b In Hospital Fall with Other Fracture Rate			0.00
•	PSI 9 Perioperative Hemorrhage or Hematoma Rate	1.59	0.00	4.15
+	PSI 10 Postoperative Acute Kidney Injury Requiring Dialysis Rate	0.76	0.00	0.00
•	PSI 11 Postoperative Respiratory Failure Rate	9.24	0.00	0.00
_	PSI 12 Perioperative Pulmonary Embolism or Deep Vein Thrombosis			
•	Rate	3.31	5.22	0.00
•	PSI 13 Postoperative Sepsis Rate	3.58	0.00	0.00
	PSI 14 Postoperative Wound Dehiscence Rate	0.83	10.42	0.00
	PSI 15 Unrecognized Abdominopelvic Accidental Puncture/Laceration Rate	1.18	0.00	0.00
<b>1</b>	SMB: Sepsis Management Bundle	56.9%	65.2%	59.4%
•	CLABSI	0.711	0.858	0.000
•	CAUTI	0.558	0.594	0.000
•	SSI COLON Surgical Site Infection	2.13	5.00	0.00
•	SSI HYST Surgical Site Infection	0.71	0.00	18.18
•	MRSA	0.047	0.000	0.000
•	CDIFF	0.671	0.287	0.000
<b>1</b>	HCOMP1A P Patients who reported that their nurses "Always" communicated well	84.6	74.6	74.4
<b>1</b>	HCOMP2A P Patients who reported that their doctors "Always" communicated well	83.3	75.2	79.1
•	HCOMP5A P Patients who reported that staff "Always" explained about medicines before giving it to them	71.0	59.9	59.5
•	HCOMP6Y P Patients who reported that YES, they were given information about what to do during their recovery at home	87.7	79.8	79.8
•	HCOMP7SA Patients who "Strongly Agree" they understood their care when they left the hospital	54.6	45.6	45.8
•	READM30HF Heart Failure 30day readmissions rate	27.7%	23.6%	17.5%
•	READM30PN Pneumonia 30day readmission rate	17.9%	20.9%	16.4%
•	Sepsis In House Mortality	10.7%	8.3%	3.9%
•	MORT30HF Heart failure 30day mortality rate	3.0%	5.2%	2.9%
•	MORT30PN Pneumonia 30day mortality rate	5.0%	3.8%	3.6%
•	Left without being seen	0.83%	2.00%	1.28%
•	Median Time from ED Arrival to Departure for Outpatients (18b)	126.00	181.3	170.1
•	Median Time from ED Arrival to Transport for Admitted Patients (ED1)	223.00	484.6	661.9

Desired Performance	Target Measures	Baseline		me Pine pital
Pe			FY23	FYTD24
•	PSI 3 Pressure Ulcer Rate	1.07	0.00	0.00
•	PSI 6 latrogenic Pneumothorax Rate	0.25	0.00	0.00
•	PSI 8a In Hospital Fall with Hip Fracture Rate	0.06	0.00	0.00
•	PSI 8b In Hospital Fall with Other Fracture Rate			0.00
	PSI 9 Perioperative Hemorrhage or Hematoma Rate	1.59	0.00	0.00
+	PSI 10 Postoperative Acute Kidney Injury Requiring Dialysis Rate	0.76		
•	PSI 11 Postoperative Respiratory Failure Rate	9.24		
	PSI 12 Perioperative Pulmonary Embolism or Deep Vein Thrombosis			
•	Rate	3.31	0.00	0.00
•	PSI 13 Postoperative Sepsis Rate	3.58		
•	PSI 14 Postoperative Wound Dehiscence Rate	0.83	0.00	0.00
+	PSI 15 Unrecognized Abdominopelvic Accidental Puncture/Laceration	1.18	0.00	0.00
<b>1</b>	SMB: Sepsis Management Bundle	56.9%	64.4%	60.0%
•	CLABSI	0.711	0.000	0.000
•	CAUTI	0.558	0.000	0.000
•	SSI COLON Surgical Site Infection	2.13		
<b>+</b>	SSI HYST Surgical Site Infection	0.71		
•	MRSA	0.047	0.000	0.000
•	CDIFF	0.671	0.000	0.000
•	HCOMP1A P Patients who reported that their nurses "Always" communicated well	84.6	78.8	78.4
•	HCOMP2A P Patients who reported that their doctors "Always" communicated well	83.3	75.6	75.2
<b>1</b>	HCOMP5A P Patients who reported that staff "Always" explained about medicines before giving it to them	71.0	67.7	76.0
•	HCOMP6Y P Patients who reported that YES, they were given information about what to do during their recovery at home	87.7	89.6	88.7
•	HCOMP7SA Patients who "Strongly Agree" they understood their care when they left the hospital	54.6	49.9	44.3
•	READM30HF Heart Failure 30day readmissions rate	27.7%	19.6%	15.0%
•	READM30PN Pneumonia 30day readmission rate	17.9%	14.0%	13.8%
•	Sepsis In House Mortality	10.7%	3.4%	0.0%
<b>+</b>	MORT30HF Heart failure 30day mortality rate	3.0%	0.0%	0.0%
•	MORT30PN Pneumonia 30day mortality rate	5.0%	3.1%	0.0%
•	Left without being seen	0.83%	1.86%	0.70%
•	Median Time from ED Arrival to Departure for Outpatients (18b)	126.00	144.5	138.5
•	Median Time from ED Arrival to Transport for Admitted Patients (ED1)	223.00	648.8	336.2

Desired Performance	Target Measures			ns County al Hospital	
		FY23	FYTD24		
+	PSI 3 Pressure Ulcer Rate	1.07	0.00	0.00	
+	PSI 6 latrogenic Pneumothorax Rate	0.25	0.00	0.00	
•	PSI 8a In Hospital Fall with Hip Fracture Rate	0.06	0.00	0.00	
•	PSI 8b In Hospital Fall with Other Fracture Rate	0.00	0.00	0.00	
•	PSI 9 Perioperative Hemorrhage or Hematoma Rate	1.59			
	PSI 10 Postoperative Acute Kidney Injury Requiring Dialysis Rate	0.76			
•	PSI 11 Postoperative Respiratory Failure Rate	9.24			
<u> </u>	PSI 12 Perioperative Pulmonary Embolism or Deep Vein Thrombosis				
•	Rate	3.31			
•	PSI 13 Postoperative Sepsis Rate	3.58			
•	PSI 14 Postoperative Wound Dehiscence Rate	0.83			
	PSI 15 Unrecognized Abdominopelvic Accidental Puncture/Laceration	1.18	0.00		
<b>1</b>	SMB: Sepsis Management Bundle	56.9%	63.0%	70.8%	
+	CLABSI	0.711	0.000	0.000	
•	CAUTI	0.558	0.000	0.000	
+	SSI COLON Surgical Site Infection	2.13	0.000	0.000	
+	SSI HYST Surgical Site Infection	0.71			
•	MRSA	0.047	0.000	0.000	
+	CDIFF	0.671	0.000	0.000	
<b>1</b>	HCOMP1A P Patients who reported that their nurses "Always" communicated well	84.6	81.6	83.4	
<b>1</b>	HCOMP2A P Patients who reported that their doctors "Always" communicated well	83.3	74.6	82.8	
•	HCOMP5A P Patients who reported that staff "Always" explained about medicines before giving it to them	71.0	73.8	82.4	
•	HCOMP6Y P Patients who reported that YES, they were given information about what to do during their recovery at home	87.7	91.9	82.2	
•	HCOMP7SA Patients who "Strongly Agree" they understood their care when they left the hospital	54.6	49.4	51.6	
•	READM30HF Heart Failure 30day readmissions rate	27.7%	15.8%	27.8%	
•	READM30PN Pneumonia 30day readmission rate	17.9%	10.2%	8.0%	
•	Sepsis In House Mortality	10.7%	5.3%	5.1%	
•	MORT30HF Heart failure 30day mortality rate	3.0%	0.0%	0.0%	
•	MORT30PN Pneumonia 30day mortality rate	5.0%	0.0%	3.6%	
•	Left without being seen	0.83%	0.50%	0.24%	
•	Median Time from ED Arrival to Departure for Outpatients (18b)	126.00	99.8	107.2	
•	Median Time from ED Arrival to Transport for Admitted Patients (ED1)	223.00	361.9	235.8	

Desired Performance	Target Measures	Russell C Baseline Hosp		•	
Pe			FY23	FYTD24	
•	PSI 3 Pressure Ulcer Rate	1.07	0.00	0.00	
•	PSI 6 latrogenic Pneumothorax Rate	0.25	0.00	0.00	
•	PSI 8a In Hospital Fall with Hip Fracture Rate	0.06	0.00	0.00	
<b>+</b>	PSI 8b In Hospital Fall with Other Fracture Rate			0.00	
•	PSI 9 Perioperative Hemorrhage or Hematoma Rate	1.59			
•	PSI 10 Postoperative Acute Kidney Injury Requiring Dialysis Rate	0.76			
•	PSI 11 Postoperative Respiratory Failure Rate	9.24			
•	PSI 12 Perioperative Pulmonary Embolism or Deep Vein Thrombosis Rate	3.31			
•	PSI 13 Postoperative Sepsis Rate	3.58			
•	PSI 14 Postoperative Wound Dehiscence Rate	0.83			
•	PSI 15 Unrecognized Abdominopelvic Accidental Puncture/Laceration Rate	1.18	0.00	0.00	
<b>1</b>	SMB: Sepsis Management Bundle	56.9%	66.1%	80.6%	
+	CLABSI	0.711	0.000	0.000	
•	CAUTI	0.558	1.130	0.000	
•	SSI COLON Surgical Site Infection	2.13			
•	SSI HYST Surgical Site Infection	0.71			
•	MRSA	0.047	0.000	0.000	
<b>+</b>	CDIFF	0.671	0.000	0.000	
•	HCOMP1A P Patients who reported that their nurses "Always" communicated well	84.6	86.2	79.9	
•	HCOMP2A P Patients who reported that their doctors "Always" communicated well	83.3	86.6	81.5	
<b>1</b>	HCOMP5A P Patients who reported that staff "Always" explained about medicines before giving it to them	71.0	74.8	70.3	
<b>1</b>	HCOMP6Y P Patients who reported that YES, they were given information about what to do during their recovery at home	87.7	92.6	86.0	
<b>1</b>	HCOMP7SA Patients who "Strongly Agree" they understood their care when they left the hospital	54.6	55.2	47.1	
•	READM30HF Heart Failure 30day readmissions rate	27.7%	32.1%	32.0%	
•	READM30PN Pneumonia 30day readmission rate	17.9%	14.4%	20.3%	
•	Sepsis In House Mortality	10.7%	4.7%	0.0%	
<b>+</b>	MORT30HF Heart failure 30day mortality rate	3.0%	5.4%	0.0%	
•	MORT30PN Pneumonia 30day mortality rate	5.0%	2.6%	1.2%	
•	Left without being seen	0.83%	1.70%	1.20%	
•	Median Time from ED Arrival to Departure for Outpatients (18b)	126.00	135.9	128.4	
•	Median Time from ED Arrival to Transport for Admitted Patients (ED1)	223.00	239.5	210.0	

Desired Performance	Target Measures			i County spital	
Pe			FY23	FYTD24	
•	PSI 3 Pressure Ulcer Rate	1.07	0.00	0.00	
•	PSI 6 latrogenic Pneumothorax Rate	0.25	0.00	0.00	
•	PSI 8a In Hospital Fall with Hip Fracture Rate	0.06	0.00	0.00	
•	PSI 8b In Hospital Fall with Other Fracture Rate			0.00	
<b>+</b>	PSI 9 Perioperative Hemorrhage or Hematoma Rate	1.59			
	PSI 10 Postoperative Acute Kidney Injury Requiring Dialysis Rate	0.76			
•	PSI 11 Postoperative Respiratory Failure Rate	9.24			
<u>`</u> _	PSI 12 Perioperative Pulmonary Embolism or Deep Vein Thrombosis				
•	Rate	3.31			
•	PSI 13 Postoperative Sepsis Rate	3.58			
•	PSI 14 Postoperative Wound Dehiscence Rate	0.83			
•	PSI 15 Unrecognized Abdominopelvic Accidental Puncture/Laceration	1.18	0.00	0.00	
•	SMB: Sepsis Management Bundle	56.9%	50.0%	66.7%	
<u>-</u>	CLABSI	0.711	0.000	0.000	
•	CAUTI	0.558	0.000	0.000	
+	SSI COLON Surgical Site Infection	2.13	0.000	0.000	
+	SSI HYST Surgical Site Infection	0.71			
•	MRSA	0.047	0.000	0.000	
+	CDIFF	0.671	0.000	0.000	
•	HCOMP1A P Patients who reported that their nurses "Always" communicated well	84.6	89.9	82.8	
•	HCOMP2A P Patients who reported that their doctors "Always" communicated well	83.3	87.4	81.6	
•	HCOMP5A P Patients who reported that staff "Always" explained about medicines before giving it to them	71.0	70.0	69.9	
•	HCOMP6Y P Patients who reported that YES, they were given information about what to do during their recovery at home	87.7	86.5	94.6	
•	HCOMP7SA Patients who "Strongly Agree" they understood their care when they left the hospital	54.6	61.0	51.1	
•	READM30HF Heart Failure 30day readmissions rate	27.7%	25.0%	7.7%	
•	READM30PN Pneumonia 30day readmission rate	17.9%	19.4%	18.9%	
•	Sepsis In House Mortality	10.7%	5.1%	2.1%	
•	MORT30HF Heart failure 30day mortality rate	3.0%	4.0%	0.0%	
<b>+</b>	MORT30PN Pneumonia 30day mortality rate	5.0%	4.1%	1.7%	
<b>+</b>	Left without being seen	0.83%	0.58%	0.31%	
•	Median Time from ED Arrival to Departure for Outpatients (18b)	126.00	141.8	138.6	
•	Median Time from ED Arrival to Transport for Admitted Patients (ED1)	223.00	881.8	693.2	

ance			Lee C	ounty	
Desired Performance	Target Measures	Baseline Commun	Communi	ity Hospital	
Ä			FY23	FYTD24	
<b>+</b>	PSI 3 Pressure Ulcer Rate	1.07	0.00	0.00	
•	PSI 6 latrogenic Pneumothorax Rate	0.25	0.00	0.00	
•	PSI 8a In Hospital Fall with Hip Fracture Rate	0.06	0.00	0.00	
•	PSI 8b In Hospital Fall with Other Fracture Rate			0.00	
•	PSI 9 Perioperative Hemorrhage or Hematoma Rate	1.59			
•	PSI 10 Postoperative Acute Kidney Injury Requiring Dialysis Rate	0.76			
<b>+</b>	PSI 11 Postoperative Respiratory Failure Rate	9.24			
	PSI 12 Perioperative Pulmonary Embolism or Deep Vein Thrombosis Rate	3.31			
	PSI 13 Postoperative Sepsis Rate	3.58			
•	PSI 14 Postoperative Wound Dehiscence Rate	0.83			
+	PSI 15 Unrecognized Abdominopelvic Accidental Puncture/Laceration Rate	1.18			
•	SMB: Sepsis Management Bundle	56.9%	57.9%	75.0%	
•	CLABSI	0.711	37.370	73.070	
•	CAUTI	0.558			
•	SSI COLON Surgical Site Infection	2.13			
•	SSI HYST Surgical Site Infection	0.71			
•	MRSA	0.047			
•	CDIFF	0.671			
•	HCOMP1A P Patients who reported that their nurses "Always" communicated well	84.6	81.9	91.5	
•	HCOMP2A P Patients who reported that their doctors "Always" communicated well	83.3	83.6	82.5	
•	HCOMP5A P Patients who reported that staff "Always" explained about medicines before giving it to them	71.0	51.9	68.5	
•	HCOMP6Y P Patients who reported that YES, they were given information about what to do during their recovery at home	87.7	88.2	85.2	
•	HCOMP7SA Patients who "Strongly Agree" they understood their care when they left the hospital	54.6	50.5	54.9	
•	READM30HF Heart Failure 30day readmissions rate	27.7%	20.0%	10.0%	
•	READM30PN Pneumonia 30day readmission rate	17.9%	16.1%	11.1%	
•	Sepsis In House Mortality	10.7%	7.1%	5.4%	
•	MORT30HF Heart failure 30day mortality rate	3.0%	0.0%	0.0%	
•	MORT30PN Pneumonia 30day mortality rate	5.0%	0.0%	0.0%	
•	Left without being seen	0.83%	1.09%	0.21%	
•	Median Time from ED Arrival to Departure for Outpatients (18b)	126.00	119.1	106.0	
•	Median Time from ED Arrival to Transport for Admitted Patients (ED1)	223.00	1400.3	358.0	

ed nance	T			enson	
Desired	Target Measures	Baseline	eline Communit	y Hospital	
Pe			FY23	FYTD24	
•	PSI 3 Pressure Ulcer Rate	1.07	0.00	0.00	
•	PSI 6 latrogenic Pneumothorax Rate	0.25	0.00	0.00	
•	PSI 8a In Hospital Fall with Hip Fracture Rate	0.06	0.00	0.00	
•	PSI 8b In Hospital Fall with Other Fracture Rate			0.00	
•	PSI 9 Perioperative Hemorrhage or Hematoma Rate	1.59			
•	PSI 10 Postoperative Acute Kidney Injury Requiring Dialysis Rate	0.76			
•	PSI 11 Postoperative Respiratory Failure Rate	9.24			
•	PSI 12 Perioperative Pulmonary Embolism or Deep Vein Thrombosis Rate	3.31			
	PSI 13 Postoperative Sepsis Rate	3.58			
•	PSI 14 Postoperative Wound Dehiscence Rate	0.83			
	PSI 15 Unrecognized Abdominopelvic Accidental Puncture/Laceration	0.03			
•	Rate	1.18			
•	SMB: Sepsis Management Bundle	56.9%			
•	CLABSI	0.711			
•	CAUTI	0.558			
•	SSI COLON Surgical Site Infection	2.13			
•	SSI HYST Surgical Site Infection	0.71			
•	MRSA	0.047			
•	CDIFF	0.671			
•	HCOMP1A P Patients who reported that their nurses "Always" communicated well	84.6	33.3	100.0	
•	HCOMP2A P Patients who reported that their doctors "Always" communicated well	83.3	66.7	33.3	
•	HCOMP5A P Patients who reported that staff "Always" explained about medicines before giving it to them	71.0	0.0	100.0	
•	HCOMP6Y P Patients who reported that YES, they were given information about what to do during their recovery at home	87.7			
•	HCOMP7SA Patients who "Strongly Agree" they understood their care when they left the hospital	54.6	0.0	33.3	
•	READM30HF Heart Failure 30day readmissions rate	27.7%			
•	READM30PN Pneumonia 30day readmission rate	17.9%	0.0%	0.0%	
•	Sepsis In House Mortality	10.7%	0.0%		
•	MORT30HF Heart failure 30day mortality rate	3.0%			
•	MORT30PN Pneumonia 30day mortality rate	5.0%	0.0%	0.0%	
•	Left without being seen	0.83%	0.90%	0.44%	
•	Median Time from ED Arrival to Departure for Outpatients (18b)	126.00	128.8	105.2	
•	Median Time from ED Arrival to Transport for Admitted Patients (ED1)	223.00	515.5		

Desired Performance	Target Measures			ck County spital	
Pe			FY23	FYTD24	
•	PSI 3 Pressure Ulcer Rate	1.07	0.00	0.00	
•	PSI 6 latrogenic Pneumothorax Rate	0.25	0.00	0.00	
•	PSI 8a In Hospital Fall with Hip Fracture Rate	0.06	0.00	0.00	
•	PSI 8b In Hospital Fall with Other Fracture Rate			0.00	
<b>+</b>	PSI 9 Perioperative Hemorrhage or Hematoma Rate	1.59			
	PSI 10 Postoperative Acute Kidney Injury Requiring Dialysis Rate	0.76			
•	PSI 11 Postoperative Respiratory Failure Rate	9.24			
<u>`</u> _	PSI 12 Perioperative Pulmonary Embolism or Deep Vein Thrombosis				
•	Rate	3.31			
•	PSI 13 Postoperative Sepsis Rate	3.58			
•	PSI 14 Postoperative Wound Dehiscence Rate	0.83			
•	PSI 15 Unrecognized Abdominopelvic Accidental Puncture/Laceration	1.18			
<b>1</b>	SMB: Sepsis Management Bundle	56.9%	75.0%	85.7%	
<u>-</u>	CLABSI	0.711	70.070	551775	
•	CAUTI	0.558			
+	SSI COLON Surgical Site Infection	2.13			
+	SSI HYST Surgical Site Infection	0.71			
•	MRSA	0.047			
•	CDIFF	0.671			
•	HCOMP1A P Patients who reported that their nurses "Always" communicated well	84.6	72.2	84.3	
<b>1</b>	HCOMP2A P Patients who reported that their doctors "Always" communicated well	83.3	83.5	79.2	
<b>1</b>	HCOMP5A P Patients who reported that staff "Always" explained about medicines before giving it to them	71.0	53.6	78.0	
•	HCOMP6Y P Patients who reported that YES, they were given information about what to do during their recovery at home	87.7	80.6	93.4	
<b>1</b>	HCOMP7SA Patients who "Strongly Agree" they understood their care when they left the hospital	54.6	38.4	78.3	
•	READM30HF Heart Failure 30day readmissions rate	27.7%	0.0%		
•	READM30PN Pneumonia 30day readmission rate	17.9%	18.8%	0.0%	
•	Sepsis In House Mortality	10.7%	0.0%	0.0%	
•	MORT30HF Heart failure 30day mortality rate	3.0%	0.0%	0.0%	
•	MORT30PN Pneumonia 30day mortality rate	5.0%	0.0%	0.0%	
•	Left without being seen	0.83%	0.08%	0.12%	
•	Median Time from ED Arrival to Departure for Outpatients (18b)	126.00	115.4	122.0	
•	Median Time from ED Arrival to Transport for Admitted Patients (ED1)	223.00	309.0	217.8	

Desired Performance	Target Measures	Johnson Baseline Community		•	
D Perf			FY23	FYTD24	
<b>+</b>	PSI 3 Pressure Ulcer Rate	1.07	0.00	0.00	
•	PSI 6 latrogenic Pneumothorax Rate	0.25	0.00	0.00	
•	PSI 8a In Hospital Fall with Hip Fracture Rate	0.06	0.00	0.00	
•	PSI 8b In Hospital Fall with Other Fracture Rate	0.00	0.00	0.00	
•	PSI 9 Perioperative Hemorrhage or Hematoma Rate	1.59		0.00	
	PSI 10 Postoperative Acute Kidney Injury Requiring Dialysis Rate	0.76			
1	PSI 11 Postoperative Respiratory Failure Rate	9.24			
•	PSI 12 Perioperative Respiratory Fundre Nate  PSI 12 Perioperative Pulmonary Embolism or Deep Vein Thrombosis				
•	Rate	3.31			
+	PSI 13 Postoperative Sepsis Rate	3.58			
•	PSI 14 Postoperative Wound Dehiscence Rate	0.83			
	PSI 15 Unrecognized Abdominopelvic Accidental Puncture/Laceration Rate	1.18			
•	SMB: Sepsis Management Bundle	56.9%	100.0%		
<u> </u>	CLABSI	0.711	100.070		
•	CAUTI	0.558			
•	SSI COLON Surgical Site Infection	2.13			
<b>I</b>	SSI HYST Surgical Site Infection	0.71			
•	MRSA	0.047			
•	CDIFF	0.671			
<b>1</b>	HCOMP1A P Patients who reported that their nurses "Always" communicated well	84.6	91.7	100.0	
•	HCOMP2A P Patients who reported that their doctors "Always" communicated well	83.3	87.5	100.0	
•	HCOMP5A P Patients who reported that staff "Always" explained about medicines before giving it to them	71.0	75.0	66.7	
•	HCOMP6Y P Patients who reported that YES, they were given information about what to do during their recovery at home	87.7	100.0	66.7	
•	HCOMP7SA Patients who "Strongly Agree" they understood their care when they left the hospital	54.6	54.2	44.4	
•	READM30HF Heart Failure 30day readmissions rate	27.7%		0.0%	
•	READM30PN Pneumonia 30day readmission rate	17.9%	20.0%	0.0%	
•	Sepsis In House Mortality	10.7%	0.0%	0.0%	
•	MORT30HF Heart failure 30day mortality rate	3.0%		0.0%	
•	MORT30PN Pneumonia 30day mortality rate	5.0%	0.0%	0.0%	
•	Left without being seen	0.83%	1.18%	0.85%	
•	Median Time from ED Arrival to Departure for Outpatients (18b)	126.00	128.9	139.1	
•	Median Time from ED Arrival to Transport for Admitted Patients (ED1)	223.00	1673.0	770.0	